

**Custom Mix Feed for Beef and Dairy Calves
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains ____ grams per ton of Amprolium.

Use Level	Indications for Use
113.5 -11,350 g/ton	As an aid in the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Top dress or mix in the daily ration. Feed ____ lbs/100 lb bodyweight/day to provide 5 mg/kg (227 mg/100#) bodyweight per day for 21 days.
113.5- 11,340 g/ton	As an aid in treatment of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> . Top dress or mix in the daily ration. Feed ____ lbs/100 lb bodyweight/day to provide 10 mg/kg (454 mg/100#) bodyweight per day for 5 days.

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use Amprolium in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Feed for Beef and Dairy Calves
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains ____ grams per ton of Amprolium.

Use Level	Indications for Use
113.5 -11,350 g/ton	As an aid in the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Top dress or mix in the daily ration. Feed ____ lbs/100 lb bodyweight/day to provide 5 mg/kg (227 mg/100#) bodyweight per day for 21 days.
113.5- 11,340 g/ton	As an aid in treatment of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> . Top dress or mix in the daily ration. Feed ____ lbs/100 lb bodyweight/day to provide 10 mg/kg (454 mg/100#) bodyweight per day for 5 days.

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use Amprolium in feeds containing bentonite.

UK Regulatory Services – 01/17