

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: BAMBERMYCINS

This feed contains ____ grams per ton of Bambermycins.

Check Use Level & Indications for Use:

<p>1 to 4 g/ton.</p> <p>Feed continuously in Type C feed at a rate of 10 -20 mg Bambermycins/head/day</p>	<p>For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter.</p>
<p>2 to 80 g/ton.</p> <p>Limitations for Use: Feed continuously at a rate of 10 – 40 mg/head/day in at least 1 lb. and not more than 10 lbs. of feed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>
<p>10-40 mg/head/day</p> <p>Limitations for Use: Free choice supplements must be formulated to provide 10-40 mg/head/day Bambermycins. Each use in a free-choice Type C medicated feed must be the subject of an approved new animal drug application (NADA) or supplemental NADA as required by 21CFR 510.455. Feed mills that use a proprietary formula are required to be licensed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>

Limitations for Type B Liquid Feed: Expiration date – 8 weeks after the date of manufacture, pH 3.8 to 7.5 and moisture content of 30 to 45%.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: BAMBERMYCINS

This feed contains ____ grams per ton of Bambermycins.

Check Use Level & Indications for Use:

<p>1 to 4 g/ton.</p> <p>Feed continuously in Type C feed at a rate of 10 -20 mg Bambermycins/head/day</p>	<p>For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter.</p>
<p>2 to 80 g/ton.</p> <p>Limitations for Use: Feed continuously at a rate of 10 – 40 mg/head/day in at least 1 lb. and not more than 10 lbs. of feed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>
<p>10-40 mg/head/day</p> <p>Limitations for Use: Free choice supplements must be formulated to provide 10-40 mg/head/day Bambermycins. Each use in a free-choice Type C medicated feed must be the subject of an approved new animal drug application (NADA) or supplemental NADA as required by 21CFR 510.455. Feed mills that use a proprietary formula are required to be licensed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>

Limitations for Type B Liquid Feed: Expiration date – 8 weeks after the date of manufacture, pH 3.8 to 7.5 and moisture content of 30 to 45%.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: BAMBERMYCINS

This feed contains ____ grams per ton of Bambermycins.

Check Use Level & Indications for Use:

<p>1 to 4 g/ton.</p> <p>Feed continuously in Type C feed at a rate of 10 -20 mg Bambermycins/head/day</p>	<p>For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter.</p>
<p>2 to 80 g/ton.</p> <p>Limitations for Use: Feed continuously at a rate of 10 – 40 mg/head/day in at least 1 lb. and not more than 10 lbs. of feed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>
<p>10-40 mg/head/day</p> <p>Limitations for Use: Free choice supplements must be formulated to provide 10-40 mg/head/day Bambermycins. Each use in a free-choice Type C medicated feed must be the subject of an approved new animal drug application (NADA) or supplemental NADA as required by 21CFR 510.455. Feed mills that use a proprietary formula are required to be licensed. Daily Bambermycins intake in excess of 20 mg/head/day have not shown to be more effective than 20 mg/head/day.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle; dairy and beef replacement heifers).</p>

Limitations for Type B Liquid Feed: Expiration date – 8 weeks after the date of manufacture, pH 3.8 to 7.5 and moisture content of 30 to 45%.