

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredient: DECOQUINATE**

This feed contains \_\_\_\_ grams per ton Decoquinatate.  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

**Check Use Level & Indications for Use:**

<p><b>12.9 – 90.8 g/ton.</b>  For Cattle (including ruminating and non-ruminating calves and veal calves)</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>.  <b>LIMITATIONS FOR USE:</b> Feed Type C feed or milk replacer at a rate to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.  ____ lbs of this feed provides 22.7 mg of Decoquinatate.</p>
<p><b>90.9 – 535.7 g/ton</b>  For Cattle (including ruminating and non-ruminating calves and veal calves)</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>.  <b>LIMITATIONS FOR USE:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration at a rate to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.  ____ lbs of this feed provides 22.7 mg of Decoquinatate.</p>

**WARNING:** Do not feed to cows producing milk for human consumption.

**Caution:** Do not use Decoquinatate in feeds containing bentonite.

**Limitations For Use:** Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinatate. Type C cattle feeds may be manufactured from Decoquinatate liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredient: DECOQUINATE**

This feed contains \_\_\_\_ grams per ton Decoquinatate.  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

**Check Use Level & Indications for Use:**

<p><b>12.9 – 90.8 g/ton.</b>  For Cattle (including ruminating and non-ruminating calves and veal calves)</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>.  <b>LIMITATIONS FOR USE:</b> Feed Type C feed or milk replacer at a rate to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.  ____ lbs of this feed provides 22.7 mg of Decoquinatate.</p>
<p><b>90.9 – 535.7 g/ton</b>  For Cattle (including ruminating and non-ruminating calves and veal calves)</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>.  <b>LIMITATIONS FOR USE:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration at a rate to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.  ____ lbs of this feed provides 22.7 mg of Decoquinatate.</p>

**WARNING:** Do not feed to cows producing milk for human consumption.

**Caution:** Do not use Decoquinatate in feeds containing bentonite.

**Limitations For Use:** Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinatate. Type C cattle feeds may be manufactured from Decoquinatate liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.