

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredient: LASALOCID**

This feed contains \_\_\_\_\_ grams per ton Lasalocid.  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p><b>10 to 30 g/ton (0.0011-0.0033%)</b></p> <p>Feed continuously in complete feed (Type C) to provide not less than 100 mg nor more than 360 mg/head/day.</p>	<p>For improved feed efficiency in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 100 to 360 mg Lasalocid/head/day.</p>
<p><b>25 to 30 g/ton (0.0027-0.0033%)</b></p> <p>Feed continuously in complete feed (Type C) to provide not less than 250 mg nor more than 360 mg/head/day.</p>	<p>For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 250 to 360 mg Lasalocid/head/day.</p>
<p><b>60 to 300 mg/head/day.</b></p> <p>Feed continuously at a rate of no less than 60 mg nor more than 300 mg/head/day. <b>Hand fed:</b> The medication must be contained in at least 1 lb. of feed. <b>Free choice:</b> The feed must be manufactured under a feed mill license utilizing an FDA-approved formulation, unless manufactured according to formulas published in 21 CFR 558.311 (e)(2),(3),or(4).</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). Intakes of Lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day.  Feed _____ to _____ lb./head/day to provide 60 to 300 mg Lasalocid/head/day.</p>
<p><b>1 mg/2.2 lb. (1 kg) of body weight per day.</b></p> <p>Hand feed to cattle weighing up to 800 lb. with a maximum of 360 mg/head/day.</p>	<p>Control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle.  Feed at the rate of _____ lb. per 100 lb. of bodyweight/day to provide 1 mg Lasalocid/2.2 lb bodyweight /day</p>

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution:** The safety of Lasalocid in unapproved species has not been established, do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A Articles or Type B feeds (cattle and sheep): Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in excess Lasalocid concentration that could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration that could be fatal to cattle and sheep.

**Special Considerations:** Type C cattle feeds may be manufactured from Lasalocid liquid Type B feeds which have a pH of 4.0 to 8.0 and bear appropriate mixing directions as follows: (1) For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. (2) For liquid Type B feeds stored in mechanical, air, or other agitation type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredient: LASALOCID**

This feed contains \_\_\_\_\_ grams per ton Lasalocid.  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p><b>10 to 30 g/ton (0.0011-0.0033%)</b></p> <p>Feed continuously in complete feed (Type C) to provide not less than 100 mg nor more than 360 mg/head/day.</p>	<p>For improved feed efficiency in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 100 to 360 mg Lasalocid/head/day.</p>
<p><b>25 to 30 g/ton (0.0027-0.0033%)</b></p> <p>Feed continuously in complete feed (Type C) to provide not less than 250 mg nor more than 360 mg/head/day.</p>	<p>For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 250 to 360 mg Lasalocid/head/day.</p>
<p><b>60 to 300 mg/head/day.</b></p> <p>Feed continuously at a rate of no less than 60 mg nor more than 300 mg/head/day. <b>Hand fed:</b> The medication must be contained in at least 1 lb. of feed. <b>Free choice:</b> The feed must be manufactured under a feed mill license utilizing an FDA-approved formulation, unless manufactured according to formulas published in 21 CFR 558.311 (e)(2),(3),or(4).</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). Intakes of Lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day.  Feed _____ to _____ lb./head/day to provide 60 to 300 mg Lasalocid/head/day.</p>
<p><b>1 mg/2.2 lb. (1 kg) of body weight per day.</b></p> <p>Hand feed to cattle weighing up to 800 lb. with a maximum of 360 mg/head/day.</p>	<p>Control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle.  Feed at the rate of _____ lb. per 100 lb. of bodyweight/day to provide 1 mg Lasalocid/2.2 lb bodyweight /day</p>

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution:** The safety of Lasalocid in unapproved species has not been established, do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A Articles or Type B feeds (cattle and sheep): Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in excess Lasalocid concentration that could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration that could be fatal to cattle and sheep.

**Special Considerations:** Type C cattle feeds may be manufactured from Lasalocid liquid Type B feeds which have a pH of 4.0 to 8.0 and bear appropriate mixing directions as follows: (1) For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. (2) For liquid Type B feeds stored in mechanical, air, or other agitation type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.