

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: POLOXALENE

This feed contains _____ grams per ton Poloxalene.

Use Level	Indications for use
1.0-2.0 g/100 lb bodyweight /day	Prevention of legume (alfalfa, clover) and wheat pasture bloat when fed continuously during exposure to bloat-producing conditions. Feed _____ to _____ lbs/100 lbs bodyweight/day to provide 1.0 to 2.0 g. poloxalene/100 lbs bodyweight/day

UK Regulatory Services – 01/17

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: POLOXALENE

This feed contains _____ grams per ton Poloxalene.

Use Level	Indications for use
1.0-2.0 g/100 lb bodyweight /day	Prevention of legume (alfalfa, clover) and wheat pasture bloat when fed continuously during exposure to bloat-producing conditions. Feed _____ to _____ lbs/100 lbs bodyweight/day to provide 1.0 to 2.0 g. poloxalene/100 lbs bodyweight/day

UK Regulatory Services – 01/17

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: POLOXALENE

This feed contains _____ grams per ton Poloxalene.

Use Level	Indications for use
1.0-2.0 g/100 lb bodyweight /day	Prevention of legume (alfalfa, clover) and wheat pasture bloat when fed continuously during exposure to bloat-producing conditions. Feed _____ to _____ lbs/100 lbs bodyweight/day to provide 1.0 to 2.0 g. poloxalene/100 lbs bodyweight/day

UK Regulatory Services – 01/17

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: POLOXALENE

This feed contains _____ grams per ton Poloxalene.

Use Level	Indications for use
1.0-2.0 g/100 lb bodyweight /day	Prevention of legume (alfalfa, clover) and wheat pasture bloat when fed continuously during exposure to bloat-producing conditions. Feed _____ to _____ lbs/100 lbs bodyweight/day to provide 1.0 to 2.0 g. poloxalene/100 lbs bodyweight/day

UK Regulatory Services – 01/17