

**Custom Mix Chicken Feed  
MEDICATED**

**Active Drug Ingredient: MONENSIN**

**This feed contains \_\_\_\_ grams per ton of MONENSIN.**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

	Continuously as sole ration 90-110 g/ton	<p><b>Chickens (broilers)</b> As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i>, <i>Eimeria tenella</i>, <i>Eimeria acervulina</i>, <i>Eimeria brunetti</i>, <i>Eimeria mivati</i> and <i>Eimeria maxima</i>.</p> <p><b>WARNING: Do not feed to laying chickens.</b> <b>CAUTION: Do not allow horses or other equines access to formulation containing Monensin.</b> <b>Ingestion of Monensin by equines has been fatal.</b> <b>In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.</b></p>
	90-110 g/ton	<p><b>Chickens (replacements, intended for use as caged layers)</b> As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i>, <i>Eimeria tenella</i>, <i>Eimeria acervulina</i>, <i>Eimeria brunetti</i>, <i>Eimeria mivati</i> and <i>Eimeria maxima</i>.</p> <p><b>WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.</b> <b>CAUTION: Do not allow horses or other equines access to formulation containing Monensin.</b> <b>Ingestion of Monensin by equines has been fatal.</b> <b>In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.</b></p>

**Custom Mix Chicken Feed  
MEDICATED**

**Active Drug Ingredient: MONENSIN**

**This feed contains \_\_\_\_ grams per ton of MONENSIN.**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

	Continuously as sole ration 90-110 g/ton	<p><b>Chickens (broilers)</b> As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i>, <i>Eimeria tenella</i>, <i>Eimeria acervulina</i>, <i>Eimeria brunetti</i>, <i>Eimeria mivati</i> and <i>Eimeria maxima</i>.</p> <p><b>WARNING: Do not feed to laying chickens.</b> <b>CAUTION: Do not allow horses or other equines access to formulation containing Monensin.</b> <b>Ingestion of Monensin by equines has been fatal.</b> <b>In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.</b></p>
	90-110 g/ton	<p><b>Chickens (replacements, intended for use as caged layers)</b> As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i>, <i>Eimeria tenella</i>, <i>Eimeria acervulina</i>, <i>Eimeria brunetti</i>, <i>Eimeria mivati</i> and <i>Eimeria maxima</i>.</p> <p><b>WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.</b> <b>CAUTION: Do not allow horses or other equines access to formulation containing Monensin.</b> <b>Ingestion of Monensin by equines has been fatal.</b> <b>In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.</b></p>