

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: ZOALENE

This feed contains ____ grams per ton of Zoalene.

Check Use Level & Indications for Use:

113.5 g/ton (0.0125%)	Chickens (Broilers) As an aid in prevention and control of coccidiosis.
113.5 g/ton (0.0125%)	Chickens (Broilers) As an aid in development of active immunity to coccidiosis. Starter ration-severe exposure to coccidiosis.
75.4-113.5 g/ton (0.0083%-0.0125%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Grower ration-severe exposure to coccidiosis.
75.4-113.5 g/ton (0.0083%-0.0125%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Starter ration-light to moderate exposure to coccidiosis.
36.3-75.4 g/ton (0.004-0.0083%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Grower ration-light to moderate exposure to coccidiosis.

CAUTION: Do not feed to laying birds; consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two-day period.

LIMITATIONS FOR USE: Grower ration not to be fed to birds over 14 weeks of age. Feed continuously as sole ration.

NOTE: Exposure conditions should be noted on label of feeds for replacement chickens.

UK Regulatory Services – 01/17

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: ZOALENE

This feed contains ____ grams per ton of Zoalene.

Check Use Level & Indications for Use:

113.5 g/ton (0.0125%)	Chickens (Broilers) As an aid in prevention and control of coccidiosis.
113.5 g/ton (0.0125%)	Chickens (Broilers) As an aid in development of active immunity to coccidiosis. Starter ration-severe exposure to coccidiosis.
75.4-113.5 g/ton (0.0083%-0.0125%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Grower ration-severe exposure to coccidiosis.
75.4-113.5 g/ton (0.0083%-0.0125%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Starter ration-light to moderate exposure to coccidiosis.
36.3-75.4 g/ton (0.004-0.0083%)	Chickens (Replacements) As an aid in development of active immunity to coccidiosis. Grower ration-light to moderate exposure to coccidiosis.

CAUTION: Do not feed to laying birds; consult a veterinarian or poultry pathologist if losses exceed 0.5% in a two-day period.

LIMITATIONS FOR USE: Grower ration not to be fed to birds over 14 weeks of age. Feed continuously as sole ration.

NOTE: Exposure conditions should be noted on label of feeds for replacement chickens.

UK Regulatory Services – 01/17