

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: DECOQUINATE

This feed contains ____ grams per ton Decoquinatate.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

12.9 – 90.8 g/ton.	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as Type C feed or milk replacer to provide 22.7 mg decoquinatate /100 lbs (0.5 mg/kg) of bodyweight /day.
90.9 – 535.7 g/ton	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as a top dress or mix into the daily ration to provide 22.7 mg decoquinatate/100 lb. (0.5 mg/kg) of bodyweight/day.

WARNING: Do not feed to goats producing milk for food.
Caution: Do not use in feeds containing bentonite.
Limitations For Use: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/17

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: DECOQUINATE

This feed contains ____ grams per ton Decoquinatate.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

12.9 – 90.8 g/ton.	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as Type C feed or milk replacer to provide 22.7 mg decoquinatate /100 lbs (0.5 mg/kg) of bodyweight /day.
90.9 – 535.7 g/ton	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as a top dress or mix into the daily ration to provide 22.7 mg decoquinatate/100 lb. (0.5 mg/kg) of bodyweight/day.

WARNING: Do not feed to goats producing milk for food.
Caution: Do not use in feeds containing bentonite.
Limitations For Use: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/17

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: DECOQUINATE

This feed contains ____ grams per ton Decoquinatate.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

12.9 – 90.8 g/ton.	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as Type C feed or milk replacer to provide 22.7 mg decoquinatate /100 lbs (0.5 mg/kg) of bodyweight /day.
90.9 – 535.7 g/ton	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as a top dress or mix into the daily ration to provide 22.7 mg decoquinatate/100 lb. (0.5 mg/kg) of bodyweight/day.

WARNING: Do not feed to goats producing milk for food.
Caution: Do not use in feeds containing bentonite.
Limitations For Use: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/17

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: DECOQUINATE

This feed contains ____ grams per ton Decoquinatate.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

12.9 – 90.8 g/ton.	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as Type C feed or milk replacer to provide 22.7 mg decoquinatate /100 lbs (0.5 mg/kg) of bodyweight /day.
90.9 – 535.7 g/ton	For the prevention of coccidiosis in young goats caused by <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> . Feed ____ lb. per 100 lb of bodyweight/day as a top dress or mix into the daily ration to provide 22.7 mg decoquinatate/100 lb. (0.5 mg/kg) of bodyweight/day.

WARNING: Do not feed to goats producing milk for food.
Caution: Do not use in feeds containing bentonite.
Limitations For Use: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/17