

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 20 grams per ton monensin.

Use Level	Indications for Use
20 g/ton. * *90%DM basis (feed continuously as sole ration)	For the prevention of coccidiosis caused by <i>Eimeria crandallis</i> , <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> .

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

LIMITATIONS FOR USE: Liquid Type B feeds must be maintained at pH 4.3-7.1. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily, even when not in use. Inadequate mixing of Monensin liquid Type B or Type C medicated feeds has resulted in increased Monensin concentration, which has been fatal to cattle and could be fatal to goats. Liquid Type B feeds shall bear an expiration date of 8 weeks after its date of manufacture.

UK Regulatory Services – 01/17

**Custom Mix Goat Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 20 grams per ton monensin.

Use Level	Indications for Use
20 g/ton. * *90%DM basis (feed continuously as sole ration)	For the prevention of coccidiosis caused by <i>Eimeria crandallis</i> , <i>Eimeria christenseni</i> and <i>Eimeria ninakohlyakimovae</i> .

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

LIMITATIONS FOR USE: Liquid Type B feeds must be maintained at pH 4.3-7.1. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily, even when not in use. Inadequate mixing of Monensin liquid Type B or Type C medicated feeds has resulted in increased Monensin concentration, which has been fatal to cattle and could be fatal to goats. Liquid Type B feeds shall bear an expiration date of 8 weeks after its date of manufacture.

UK Regulatory Services – 01/17