

**Custom Mix Feed
MEDICATED**

Active Drug Ingredient: Rabon

This feed contains ____ g/ton Rabon

Check Use Level & Indication for Use

0.00015 lb. per 100 lb. bodyweight/day (0.07 g/cwt./day)	Cattle (beef and dairy) Control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle. Feed at the rate of ____lb./100 lb. bodyweight/day to provide the correct amount of Rabon.
0.00015 lb per 100 lb. bodyweight/day (0.07 g/cwt./day)	Horses Control of fecal flies in manure of treated horses. Prevents development of house flies and stable flies in the manure of treated horses. Feed at the rate of ____lb./100 lb. bodyweight/day to provide the correct amount of Rabon.
0.00010 lb per 100 lb. bodyweight/day (0.05 g/cwt./day)	Swine Control of fecal flies in manure of treated swine. Prevents development of house flies in the manure of treated swine. Feed at the rate of ____lb./100 lb. bodyweight/day to provide the correct amount of Rabon.

Caution:

Keep out of reach of children. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. For personal protective equipment (PPE), mixers and handlers must wear: long-sleeved shirt and pants, shoes and socks and chemical resistant gloves. User safety requirements are to follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. User safety recommendations are that users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash water.

Note: Rabon may be used in feed containing any approved drug unless contraindicated by the drug approval, e.g. other cholinesterase inhibitors.

Please Note:

FDA STATUS: Product is a food additive (40 CFR 186.950-formerly 21 CFR 561.91). When used in medicated feeds, medicated feed application requirement is determined by regulatory status of the drug.

EPA STATUS: Product is a pesticide when used in non-medicated feeds. EPA registration is required for feeds offered for sale except when custom blended per the provisions of 40 CFR 167.3. In medicated feeds, the product is a food additive and no EPA registration is required when the source of Rabon is an EPA-registered product.