

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

**Custom Mix Pheasant Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
0.0175% (159 g/ton) Feed continuously as the sole ration.	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i>, <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i>.

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17