

**Custom Mix Poultry Feed
MEDICATED**

Active Drug Ingredient: AMPROLIUM

This feed contains ____ grams per ton of AMPROLIUM.
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

Continuously 113.5-227 g/ton (0.0125-0.025%)	Chickens (broilers) For prevention of coccidiosis where immunity to coccidiosis is not desired.
72.6-113.5 g/ton (0.008-0.0125%)	Chickens (broilers) For prevention of coccidiosis caused by <i>Eimeria tenella</i> only.
Continuously 113.5-227 g/ton (0.0125-0.025%)	Chickens (replacements) For prevention of coccidiosis where immunity to coccidiosis is not desired.
113.5 g/ton (0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of severe exposure to coccidiosis up to 5 weeks of age.
72.6-113.5 g/ton (0.008-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of severe exposure to coccidiosis from 5-8 weeks of age.
36.3-113.5 g/ton (0.004-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of severe exposure to coccidiosis over 8 weeks of age.
72.6-113.5 g/ton (0.008-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of moderate exposure to coccidiosis up to 5 weeks of age.
54.5-113.5 g/ton (0.006-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of moderate exposure to coccidiosis from 5-8 weeks of age.
36.3-113.5 g/ton (0.004-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of moderate exposure to coccidiosis over 8 weeks of age.
36.3-113.5 g/ton (0.004-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of slight exposure to coccidiosis up to 5 weeks of age.
36.3-113.5 g/ton (0.004-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of slight exposure to coccidiosis from 5-8 weeks of age.

Check Use Level & Indications for Use:

36.3-113.5 g/ton (0.004-0.0125%)	Chickens (replacements) For development of active immunity to coccidiosis under conditions of slight exposure to coccidiosis over 8 weeks of age.
Continuously 113.5 g/ton	Laying Hens For prevention of coccidiosis.
113.5 g/ton (0.0125%) Treat for 2 weeks	Laying Hens For treatment of coccidiosis in moderate outbreaks.
227 g/ton (0.025%) Treat for 2 weeks	Laying Hens For treatment of coccidiosis in severe outbreaks.
113.5-227 g/ton (0.0125-0.025%)	Turkeys For prevention of coccidiosis. Limitation for use: Feed continuously as the sole ration.

WARNING: Use as a sole source of Amprolium.

SPECIAL CONSIDERATION: Do not use Amprolium in feeds containing bentonite.