

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17

**Custom Mix Bobwhite Quail Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains 73 grams per ton of Monensin.

Use Level	Indications for Use
73 g/ton (as is) WARNING: Feed continuously as the sole ration.	For the prevention of coccidiosis in growing Bobwhite quail caused by <i>Eimeria dispersa</i> and <i>Eimeria lettyae</i> .

Caution: Do not allow horses or other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by equines has been fatal.

UK Regulatory Services – 01/17