

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient: DECOQUINATE**

**This feed contains \_\_\_\_\_ grams per ton decoquinatate.**  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level	Indications for Use
<p><b>For Young Sheep</b>  12.9 – 90.8 g/ton.</p> <p><b>Limitations:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.</p>	<p>For the prevention of coccidiosis in young sheep caused by <i>Eimeria bakuensis</i>, <i>Eimeria crandallis</i>, <i>Eimeria ovinoidalis</i> and <i>Eimeria parva</i>.</p> <p><b>Limitations:</b> Feed Type C medicated feed or milk replacer at a rate of _____ lb./100 lb. of body weight /day to provide 22.7 mg/100 lb (0.5 mg/kg) of bodyweight per day.</p>
<p><b>For Young Sheep</b>  90.9 – 535.7 g/ton.</p> <p><b>Limitations:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration to provide 22.7 mg/100 lb (0.5 mg/kg) of bodyweight/day.</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bakuensis</i>, <i>Eimeria crandallis</i>, <i>Eimeria ovinoidialis</i> and <i>Eimeria parva</i>.</p> <p>Feed _____ lb./100 lb. of bodyweight/day to provide 22.7 mg/100 lb. (0.5mg/kg) of bodyweight/day.</p>

**WARNING:** Do not feed to sheep producing milk for human consumption.

**CAUTION:** Do not use Decoquinatate in feeds containing bentonite.

**LIMITATIONS FOR USE:** Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient: DECOQUINATE**

**This feed contains \_\_\_\_\_ grams per ton decoquinatate.**  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level	Indications for Use
<p><b>For Young Sheep</b>  12.9 – 90.8 g/ton.</p> <p><b>Limitations:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration to provide 22.7 mg per 100 lb. of bodyweight (0.5 mg/kg) per day.</p>	<p>For the prevention of coccidiosis in young sheep caused by <i>Eimeria bakuensis</i>, <i>Eimeria crandallis</i>, <i>Eimeria ovinoidalis</i> and <i>Eimeria parva</i>.</p> <p><b>Limitations:</b> Feed Type C medicated feed or milk replacer at a rate of _____ lb./100 lb. of body weight /day to provide 22.7 mg/100 lb (0.5 mg/kg) of bodyweight per day.</p>
<p><b>For Young Sheep</b>  90.9 – 535.7 g/ton.</p> <p><b>Limitations:</b> Feed Type C medicated feed supplements as a top dress or mix into the daily ration to provide 22.7 mg/100 lb (0.5 mg/kg) of bodyweight/day.</p>	<p>For the prevention of coccidiosis caused by <i>Eimeria bakuensis</i>, <i>Eimeria crandallis</i>, <i>Eimeria ovinoidialis</i> and <i>Eimeria parva</i>.</p> <p>Feed _____ lb./100 lb. of bodyweight/day to provide 22.7 mg/100 lb. (0.5mg/kg) of bodyweight/day.</p>

**WARNING:** Do not feed to sheep producing milk for human consumption.

**CAUTION:** Do not use Decoquinatate in feeds containing bentonite.

**LIMITATIONS FOR USE:** Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.