

**Custom Mix Swine Feed
MEDICATED**

Active Drug Ingredient: **BACITRACIN
METHYLENE DISALICYLATE (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

| | |
|---|--|
| 10 to 30 g/ton. For growing and finishing swine. | For increased rate of weight gain and improved feed efficiency. |
| 250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. Not for use in swine weighing more than 250 lb. | For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition. |
| 250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. | For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. |

**Custom Mix Swine Feed
MEDICATED**

Active Drug Ingredient: **BACITRACIN
METHYLENE DISALICYLATE (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

| | |
|---|--|
| 10 to 30 g/ton. For growing and finishing swine. | For increased rate of weight gain and improved feed efficiency. |
| 250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. Not for use in swine weighing more than 250 lb. | For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition. |
| 250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. | For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. |

**Custom Mix Swine Feed
MEDICATED**

Active Drug Ingredient: **BACITRACIN
METHYLENE DISALICYLATE (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

| | |
|---|--|
| 10 to 30 g/ton. For growing and finishing swine. | For increased rate of weight gain and improved feed efficiency. |
| 250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. Not for use in swine weighing more than 250 lb. | For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition. |
| 250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be the sole ration. | For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. |