

**CUSTOM MIX SWINE FEED
Medicated**

Active Drug Ingredients: CARBADOX

This feed contains ___ g/ton of Carbadox

Check Use Level & Indications for Use:

10-25 g/ton. (0.0011- 0.00275%)	For increase in rate of weight gain and improvement of feed efficiency
50g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>); for increased rate of weight gain and improvement of feed efficiency.

WARNING: Do not feed to swine within 42 days before slaughter.

Caution: Do not use in feeds containing bentonite. Not for use in pregnant swine or swine intended for breeding purposes.

UK Regulatory Services – 01/17

**CUSTOM MIX SWINE FEED
Medicated**

Active Drug Ingredients: CARBADOX

This feed contains ___ g/ton of Carbadox

Check Use Level & Indications for Use:

10-25 g/ton. (0.0011- 0.00275%)	For increase in rate of weight gain and improvement of feed efficiency
50g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>); for increased rate of weight gain and improvement of feed efficiency.

WARNING: Do not feed to swine within 42 days before slaughter.

Caution: Do not use in feeds containing bentonite. Not for use in pregnant swine or swine intended for breeding purposes.

UK Regulatory Services – 01/17

**CUSTOM MIX SWINE FEED
Medicated**

Active Drug Ingredients: CARBADOX

This feed contains ___ g/ton of Carbadox

Check Use Level & Indications for Use:

10-25 g/ton. (0.0011- 0.00275%)	For increase in rate of weight gain and improvement of feed efficiency
50g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>); for increased rate of weight gain and improvement of feed efficiency.

WARNING: Do not feed to swine within 42 days before slaughter.

Caution: Do not use in feeds containing bentonite. Not for use in pregnant swine or swine intended for breeding purposes.

UK Regulatory Services – 01/17

**CUSTOM MIX SWINE FEED
Medicated**

Active Drug Ingredients: CARBADOX

This feed contains ___ g/ton of Carbadox

Check Use Level & Indications for Use:

10-25 g/ton. (0.0011- 0.00275%)	For increase in rate of weight gain and improvement of feed efficiency
50g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>); for increased rate of weight gain and improvement of feed efficiency.

WARNING: Do not feed to swine within 42 days before slaughter.

Caution: Do not use in feeds containing bentonite. Not for use in pregnant swine or swine intended for breeding purposes.

UK Regulatory Services – 01/17