

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: CARBADOX and
PYRANTEL TARTRATE.**

Use Level	Indications for Use
Carbadox: 50 g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>).
Pyrantel Tartrate: 96 g/ton (0.0106%)	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections and aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Do not feed to swine weighing more than 75 lb. bodyweight. Do not feed to swine within 10 weeks of slaughter. Do not use in complete feeds containing less than 15% crude protein.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Feed continuously as the sole ration.

UK Regulatory Services – 01/17

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: CARBADOX and
PYRANTEL TARTRATE.**

Use Level	Indications for Use
Carbadox: 50 g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>).
Pyrantel Tartrate: 96 g/ton (0.0106%)	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections and aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Do not feed to swine weighing more than 75 lb. bodyweight. Do not feed to swine within 10 weeks of slaughter. Do not use in complete feeds containing less than 15% crude protein.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Feed continuously as the sole ration.

UK Regulatory Services – 01/17

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: CARBADOX and
PYRANTEL TARTRATE.**

Use Level	Indications for Use
Carbadox: 50 g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>).
Pyrantel Tartrate: 96 g/ton (0.0106%)	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections and aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Do not feed to swine weighing more than 75 lb. bodyweight. Do not feed to swine within 10 weeks of slaughter. Do not use in complete feeds containing less than 15% crude protein.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Feed continuously as the sole ration.

UK Regulatory Services – 01/17

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: CARBADOX and
PYRANTEL TARTRATE.**

Use Level	Indications for Use
Carbadox: 50 g/ton (0.0055%)	For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by <i>Salmonella choleraesuis</i>).
Pyrantel Tartrate: 96 g/ton (0.0106%)	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections and aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Do not feed to swine weighing more than 75 lb. bodyweight. Do not feed to swine within 10 weeks of slaughter. Do not use in complete feeds containing less than 15% crude protein.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Feed continuously as the sole ration.

UK Regulatory Services – 01/17