

Custom Mix Swine Feed
MEDICATED

Active Drug Ingredient: PYRANTEL
TARTRATE.

Check Use Level & Indications for Use:

96 g/ton (0.0106%) Feed continuously as the sole ration in a complete feed.	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.
96 g/ton (0.0106%) Feed for 3 days as the sole ration in a complete ration	For the removal and control of large roundworm (<i>Ascaris suum</i>) infections.
800 g/ton (0.0881%) As a single therapeutic treatment in a complete feed; feed at the rate of 1 lb. of feed per 40 lb. of bodyweight for animals up to 200 lb., and 5 lb. of feed per head for animals 200 lb. or over.	For the removal and control of large roundworm (<i>Ascaris suum</i>) and nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Withdraw 24 hours prior to slaughter.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17

Custom Mix Swine Feed
MEDICATED

Active Drug Ingredient: PYRANTEL
TARTRATE.

Check Use Level & Indications for Use:

96 g/ton (0.0106%) Feed continuously as the sole ration in a complete feed.	Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.
96 g/ton (0.0106%) Feed for 3 days as the sole ration in a complete ration	For the removal and control of large roundworm (<i>Ascaris suum</i>) infections.
800 g/ton (0.0881%) As a single therapeutic treatment in a complete feed; feed at the rate of 1 lb. of feed per 40 lb. of bodyweight for animals up to 200 lb., and 5 lb. of feed per head for animals 200 lb. or over.	For the removal and control of large roundworm (<i>Ascaris suum</i>) and nodular worm (<i>Oesophagostomum</i>) infections.

WARNING: Withdraw 24 hours prior to slaughter.

CAUTION: Consult veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

UK Regulatory Services – 01/17