

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

Calves (up to 250 lb.) 10 mg/lb. bodyweight/day*	For treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to chlortetracycline. Feed ____ lb./100 lb. bodyweight to provide 10 mg/lb. bodyweight/day of chlortetracycline. LIMITS FOR USE: In milk replacers or starter feeds.
Growing Cattle (over 400 lb.) 70 mg/head/day*	Reduction of incidence of liver abscesses. Feed ____ lb./head/day to provide 70 mg chlortetracycline.
Beef cattle 350 mg/head/day*	Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella spp.</i> susceptible to Chlortetracycline. Feed ____ lb./head/day to provide 350 mg chlortetracycline.
Beef cattle (under 700 lb.) 350 mg/head/day*	For control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. Feed ____ lb./head/day to provide 350 mg chlortetracycline.
Beef cattle (over 700 lb.) 0.5 mg/lb. bodyweight/day*	Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. Feed ____ lb. daily to provide 0.5 mg of chlortetracycline/lb. bodyweight. <i>Example: To provide 0.5 mg/lb. bodyweight at a feed intake of 0.5 lb/100 lb. bodyweight, mix a feed containing 200g Chlortetracycline</i>
Calves, beef and non-lactating dairy cattle 10 mg/lb. bodyweight/day*	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. Feed ____ lb./100 lb. bodyweight/day to provide 10 mg chlortetracycline/lb. bodyweight/day. Limitations for Use: Feed for not more than 5 days. Feed approximately 400 g/ton varying with bodyweight and feed consumption to provide 10 mg/lb. per day of chlortetracycline.
Beef and non-lactating dairy cattle 0.5-2.0 mg/lb. bodyweight/day	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. LIMITATIONS: In free-choice cattle feeds such as feed blocks or salt-mineral mixes; Free choice feed must be manufactured from NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin) using an FDA-approved formulation. Feed mill license not required if following free-choice loose-mineral formulation published in 21 CFR 558.128.
Calves, beef and non-lactating dairy cattle 500 – 4,000 g/ton	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. LIMITATIONS: Feed continuously for not more than 5 days to provide 10 mg/lb. bodyweight per day; must be manufactured under NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin).
Beef and non-lactating dairy cattle 6,000 g/ton	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline.
Calves, beef and non-lactating dairy cattle 4,000-20,000 g/ton	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. LIMITATIONS: As a top dress, varying with bodyweight and feed consumption, to provide 10 mg/lb. bodyweight per day. Treat for not more than 5 days, must be manufactured from Aureomycin Type A medicated articles manufactured under NADA 048-761 or Deracin Type A medicated articles manufactured under ANADA 200-510.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

*Withdrawal times may vary by manufacturer. Read label for specific withdrawal times.

Withdrawal time for the use checked above is _____ prior to slaughter.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.