

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredient: CHLORTETRACYCLINE**

This feed contains \_\_\_\_ grams per ton of Chlortetracycline (CTC).  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

Calves (up to 250 lb.)  <b>10 mg/lb. bodyweight/day*</b>	For treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to chlortetracycline.  <b>Feed ____ lb./100 lb. bodyweight to provide 10 mg/lb. bodyweight/day of chlortetracycline.</b> <b>LIMITS FOR USE: In milk replacers or starter feeds.</b>
Growing Cattle (over 400 lb.)  <b>70 mg/head/day*</b>	For reduction of liver condemnation due to liver abscesses.  <b>Feed ____ lb./head/day to provide 70 mg chlortetracycline.</b>
Beef cattle  <b>350 mg/head/day*</b>	Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella spp.</i> susceptible to Chlortetracycline.  <b>Feed ____ lb./head/day to provide 350 mg chlortetracycline.</b>
Beef cattle (under 700 lb.)  <b>350 mg/head/day*</b>	For control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline.  <b>Feed ____ lb./head/day to provide 350 mg chlortetracycline.</b>
Beef cattle (over 700 lb.)  <b>0.5 mg/lb. bodyweight/day*</b>	Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline.  <b>Feed ____ lb. daily to provide 0.5 mg of chlortetracycline/lb. bodyweight.</b> <i>Example: To provide 0.5 mg/lb. bodyweight at a feed intake of 0.5 lb/100 lb. bodyweight, mix a feed containing 200g Chlortetracycline</i>
Calves, beef and non-lactating dairy cattle  <b>10 mg/lb. bodyweight/day*</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.  <b>Feed ____ lb./100 lb. bodyweight/day to provide 10 mg chlortetracycline/lb. bodyweight/day.</b> <b>Limits for Use: Feed for not more than 5 days. Feed approximately 400 g/ton varying with bodyweight and feed consumption to provide 10 mg/lb. per day of chlortetracycline.</b>
Beef and non-lactating dairy cattle  <b>0.5-2.0 mg/lb. bodyweight/day</b>	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. <b>LIMITATIONS: In free-choice cattle feeds such as feed blocks or salt-mineral mixes; Free choice feed must be manufactured from NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin) using an FDA-approved formulation. Feed mill license not required if following free-choice loose-mineral formulation published in 21 CFR 558.128.</b>
Calves, beef and non-lactating dairy cattle  <b>500 – 4,000 g/ton</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. <b>LIMITATIONS: Hand feed continuously for not more than 5 days to provide 10 mg/lb. bodyweight per day; must be manufactured under NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin).</b>
Beef and non-lactating dairy cattle  <b>6,000 g/ton</b>	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline.
Calves, beef and non-lactating dairy cattle  <b>4,000-20,000 g/ton</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. <b>LIMITATIONS: As a top dress, varying with bodyweight and feed consumption, to provide 10 mg/lb. bodyweight per day. Treat for not more than 5 days, must be manufactured from Aureomycin Type A medicated articles manufactured under NADA 048-761 or Deracin Type A medicated articles manufactured under ANADA 200-510.</b>

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

\*Withdrawal times may vary by manufacturer. Read label for specific withdrawal times.

**Withdrawal time for the use checked above is \_\_\_\_\_ prior to slaughter.**

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.