

**Custom Mix Cattle Feed
Chlortetracycline / Decoquinatate
MEDICATED**

Active Drug Ingredients:

_____ grams per ton **CHLORTETRACYCLINE (CTC)** and
_____ grams per ton **DECOQUINATE**
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use:
<p>CTC 400g/ton (10mg/lb bodyweight/day)</p> <p>and</p> <p>Decoquinatate 13.6--27.2 g/ton</p>	<p>Calves, beef and nonlactating dairy cattle For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuerni</i>. For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.</p> <p>Feed _____ lbs./100 lb bodyweight/day to provide 22.7 mg Decoquinatate and 1 g CTC per 100 lb. bodyweight per day for not more than 5 days.</p> <p>Limitations for Use Feed approximately 400 g/ton of chlortetracycline varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Type C feed may be prepared from Type B feed containing 535.8 to 5,440 g/ton Decoquinatate and 6,700 to 80,000 g/ton Chlortetracycline.</p>
<p>CTC 500 – 4000 g/ton</p> <p>and</p> <p>Decoquinatate 12.9-90.8 g/ton</p>	<p>Calves, beef and nonlactating dairy cattle For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>. For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.</p> <p>Feed _____ lbs./100 lb bodyweight/day to provide 22.7 mg Decoquinatate and 1 g CTC per 100 lb. bodyweight per day for not more than 5 days.</p>
<p>CTC 4,000 to 20,000 g/ton</p> <p>and</p> <p>Decoquinatate 90.9 to 535.7 g/ton</p>	<p>Calves, beef and nonlactating dairy cattle For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>. For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.</p> <p>LIMITATIONS FOR USE: Feed Type C medicated supplements as a top dress or mix into daily ration to provide 22.7 mg Decoquinatate and 1 g CTC per 100 lb. bodyweight per day for not more than 5 days.</p>

LIMITATIONS FOR USE: Feed Type C feed at a rate to provide 22.7 mg Decoquinatate and 1 g Chlortetracycline per 100 lb. of bodyweight (0.5 mg/kg) per day for not more than 5 days. When consumed, feed 22.7 mg Decoquinatate per 100 lb. bodyweight per day for total of 28 days to prevent coccidiosis. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinatate. Type C cattle feeds may be manufactured from Decoquinatate liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.

WARNING: Do not feed to animals producing milk for food. A withdrawal period has not been established for this product in pre-ruminating calves. Do not feed to calves to be processed for veal. Withdraw 24 hours prior to slaughter when manufactured from ChlorMax Type A medicated articles and Pennchlor Type A medicated articles. Zero withdrawal time when manufactured from Aureomycin Type A medicated articles manufactured by NADA sponsor No. 054771 (Zoetis).

CAUTION: Do not use Decoquinatate in feeds containing bentonite.

CAUTION: Federal law restricts medicated feed containing this Veterinary feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.