

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. bodyweight daily*</p> <p>In starter feeds or milk replacers for calves.</p> <p>Feed continuously daily for 7 to 14 days.</p>	<p>For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline in calves (up to 250 lb.)</p> <p>Feed at the rate of _____ lb./100 lb. bodyweight/day.</p>
<p>75 mg/head/day</p> <p>For Beef cattle. Use continuously</p>	<p>For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses.</p> <p>Feed _____ lb./head/day.</p>
<p>0.5 to 2.0 g/head/day</p> <p>For beef cattle. Feed 3-5 days before and after arrival in feedlot.</p>	<p>Prevention and treatment of the early stages of shipping fever complex.</p> <p>Feed __ to __ lb./head/day.</p>
<p>10 mg/lb. bodyweight daily</p> <p>For calves including prurminating (veal) calves, beef cattle and nonlacting dairy cattle.</p> <p>Feed continuously for 7-14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.</p> <p>Feed _____ lb./100 lb. bodyweight/day.</p>

***NOTE: This product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.**

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. bodyweight daily*</p> <p>In starter feeds or milk replacers for calves.</p> <p>Feed continuously daily for 7 to 14 days.</p>	<p>For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline in calves (up to 250 lb.)</p> <p>Feed at the rate of _____ lb./100 lb. bodyweight/day.</p>
<p>75 mg/head/day</p> <p>For Beef cattle. Use continuously</p>	<p>For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses.</p> <p>Feed _____ lb./head/day.</p>
<p>0.5 to 2.0 g/head/day</p> <p>For beef cattle. Feed 3-5 days before and after arrival in feedlot.</p>	<p>Prevention and treatment of the early stages of shipping fever complex.</p> <p>Feed __ to __ lb./head/day.</p>
<p>10 mg/lb. bodyweight daily</p> <p>For calves including prurminating (veal) calves, beef cattle and nonlacting dairy cattle.</p> <p>Feed continuously for 7-14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.</p> <p>Feed _____ lb./100 lb. bodyweight/day.</p>

***NOTE: This product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.**

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.