

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. bodyweight daily</p> <p>In starter feeds or milk replacers for calves.</p> <p>Feed continuously daily for 7 to 14 days.</p>	<p>For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline in calves (up to 250 lb.)</p> <p>Feed at the rate of _____ lb./100 lb. bodyweight/day.</p>
<p>75 mg/head/day</p> <p>Use continuously</p>	<p>For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses.</p> <p>Feed _____ lb./head/day.</p>
<p>0.5 to 2.0 g/head/day</p> <p>For beef cattle. Feed 3-5 days before and after arrival in feedlot.</p>	<p>Prevention and treatment of the early stages of shipping fever complex.</p> <p>Feed __ to __ lb./head/day.</p>
<p>10 mg/lb. bodyweight daily</p> <p>For calves including pruminating (veal) calves, beef cattle and nonlacting dairy cattle.</p> <p>Feed continuously for 7-14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.</p> <p>Feed _____ lb./100 lb. bodyweight/day.</p>

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. bodyweight daily</p> <p>In starter feeds or milk replacers for calves.</p> <p>Feed continuously daily for 7 to 14 days.</p>	<p>For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline in calves (up to 250 lb.)</p> <p>Feed at the rate of _____ lb./100 lb. bodyweight/day.</p>
<p>75 mg/head/day</p> <p>Use continuously</p>	<p>For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses.</p> <p>Feed _____ lb./head/day.</p>
<p>0.5 to 2.0 g/head/day</p> <p>For beef cattle. Feed 3-5 days before and after arrival in feedlot.</p>	<p>Prevention and treatment of the early stages of shipping fever complex.</p> <p>Feed __ to __ lb./head/day.</p>
<p>10 mg/lb. bodyweight daily</p> <p>For calves including pruminating (veal) calves, beef cattle and nonlacting dairy cattle.</p> <p>Feed continuously for 7-14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.</p> <p>Feed _____ lb./100 lb. bodyweight/day.</p>

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.