

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: TYLOSIN

This feed contains ____ grams per ton tylosin.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

8 to 10 g/ton. * 90% DM basis	For reduction of incidence of liver abscesses in beef cattle caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium (Actinomyces) pyogenes.</i>
Feed continuously at a rate of 60-90 mg/head/day.	
	Feed ____ to ____ lbs/head/day continuously to provide 60-90 mg Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: TYLOSIN

This feed contains ____ grams per ton tylosin.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

8 to 10 g/ton. * 90% DM basis	For reduction of incidence of liver abscesses in beef cattle caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium (Actinomyces) pyogenes.</i>
Feed continuously at a rate of 60-90 mg/head/day.	
	Feed ____ to ____ lbs/head/day continuously to provide 60-90 mg Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: TYLOSIN

This feed contains ____ grams per ton tylosin.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

8 to 10 g/ton. * 90% DM basis	For reduction of incidence of liver abscesses in beef cattle caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium (Actinomyces) pyogenes.</i>
Feed continuously at a rate of 60-90 mg/head/day.	
	Feed ____ to ____ lbs/head/day continuously to provide 60-90 mg Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.