

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ___ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

100-200 g/ton (0.011-0.022%)	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.
200-400 g/ton (0.022-0.044%)	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.
500 g/ton (0.055%)	Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to Chlortetracycline. LIMITS FOR USE: (1)Feed for 5 days. Zero-day withdrawal time when formulated from Aureomycin Type A medicated articles or Type B medicated feeds under NADA 48-761 or Deracin Type A medicated articles or Type B medicated feeds under ANADA 200-510. (2) All other Chlortetracycline premixes: Feed for 5 days. Withdraw 24 hours prior to slaughter*. Do not feed to chickens producing eggs for human consumption.

WARNING: * Withdrawal times vary by manufacturer. Read drug premix label for specific withdrawal time and limits for use.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ___ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

100-200 g/ton (0.011-0.022%)	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.
200-400 g/ton (0.022-0.044%)	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.
500 g/ton (0.055%)	Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to Chlortetracycline. LIMITS FOR USE: (1)Feed for 5 days. Zero-day withdrawal time when formulated from Aureomycin Type A medicated articles or Type B medicated feeds under NADA 48-761 or Deracin Type A medicated articles or Type B medicated feeds under ANADA 200-510. (2) All other Chlortetracycline premixes: Feed for 5 days. Withdraw 24 hours prior to slaughter*. Do not feed to chickens producing eggs for human consumption.

WARNING: * Withdrawal times vary by manufacturer. Read drug premix label for specific withdrawal time and limits for use.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.