

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient:  
OXYTETRACYCLINE**

This feed contains \_\_\_\_ grams per ton of  
Oxytetracycline.  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b>10 mg/lb. of bodyweight daily.</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
Feed continuously for 7-14 days.	<b>WARNING: Five day withdrawal period required at this dosage.</b>

**SPECIAL CONSIDERATION:** For use in dry animal feed  
only. Not for use in liquid feeds supplements.

**CAUTION:** Federal law restricts medicated feed containing  
this Veterinary Feed Directive (VFD) drug to use by or on  
the order of a licensed veterinarian.

UK Regulatory Services – 01/17

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient:  
OXYTETRACYCLINE**

This feed contains \_\_\_\_ grams per ton of  
Oxytetracycline.  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b>10 mg/lb. of bodyweight daily.</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
Feed continuously for 7-14 days.	<b>WARNING: Five day withdrawal period required at this dosage.</b>

**SPECIAL CONSIDERATION:** For use in dry animal feed  
only. Not for use in liquid feeds supplements.

**CAUTION:** Federal law restricts medicated feed containing  
this Veterinary Feed Directive (VFD) drug to use by or on  
the order of a licensed veterinarian.

UK Regulatory Services – 01/17

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient:  
OXYTETRACYCLINE**

This feed contains \_\_\_\_ grams per ton of  
Oxytetracycline.  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b>10 mg/lb. of bodyweight daily.</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
Feed continuously for 7-14 days.	<b>WARNING: Five day withdrawal period required at this dosage.</b>

**SPECIAL CONSIDERATION:** For use in dry animal feed  
only. Not for use in liquid feeds supplements.

**CAUTION:** Federal law restricts medicated feed containing  
this Veterinary Feed Directive (VFD) drug to use by or on  
the order of a licensed veterinarian.

UK Regulatory Services – 01/17

**Custom Mix Sheep Feed  
MEDICATED**

**Active Drug Ingredient:  
OXYTETRACYCLINE**

This feed contains \_\_\_\_ grams per ton of  
Oxytetracycline.  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b>10 mg/lb. of bodyweight daily.</b>	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
Feed continuously for 7-14 days.	<b>WARNING: Five day withdrawal period required at this dosage.</b>

**SPECIAL CONSIDERATION:** For use in dry animal feed  
only. Not for use in liquid feeds supplements.

**CAUTION:** Federal law restricts medicated feed containing  
this Veterinary Feed Directive (VFD) drug to use by or on  
the order of a licensed veterinarian.

UK Regulatory Services – 01/17