

**Custom Mix Swine Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

50-100 g/ton (growing swine)	Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group <i>E Streptococci</i> susceptible to Chlortetracycline.
400 g/ton (breeding swine)	Control of leptospirosis (reducing the incidence of abortion and shedding of Leptospirae) caused by <i>Leptospira pomona</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for not more than 14 days.
10 mg/lb. bodyweight/day	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed for not more than 14 days. Feed approximately 400 g/ton, varying with bodyweight and feed consumption to provide 10 mg/lb. bodyweight per day. For products manufactured under NADA 046-699 (Chlormax), NADA 048-761 (Aureomycin), NADA 138-935 (Pennchlor) or ANADA 200-510 (Deracin), zero-day withdrawal.
10 mg/lb. bodyweight/day	For the control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed for not more than 14 days. Feed approximately 400 g/ton, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. FDA approval provides only for use of products manufactured under NADA 048-761 (Aureomycin), 046-699 (Chlormax) or ANADA 200-510 (Deracin).

WARNING: *Withdrawal times vary by manufacturer. Read drug premix label for specific withdrawal time.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Swine Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

50-100 g/ton (growing swine)	Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group <i>E Streptococci</i> susceptible to Chlortetracycline.
400 g/ton (breeding swine)	Control of leptospirosis (reducing the incidence of abortion and shedding of Leptospirae) caused by <i>Leptospira pomona</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for not more than 14 days.
10 mg/lb. bodyweight/day	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed for not more than 14 days. Feed approximately 400 g/ton, varying with bodyweight and feed consumption to provide 10 mg/lb. bodyweight per day. For products manufactured under NADA 046-699 (Chlormax), NADA 048-761 (Aureomycin), NADA 138-935 (Pennchlor) or ANADA 200-510 (Deracin), zero-day withdrawal.
10 mg/lb. bodyweight/day	For the control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed for not more than 14 days. Feed approximately 400 g/ton, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. FDA approval provides only for use of products manufactured under NADA 048-761 (Aureomycin), 046-699 (Chlormax) or ANADA 200-510 (Deracin).

WARNING: *Withdrawal times vary by manufacturer. Read drug premix label for specific withdrawal time.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.