

**Custom Mix Swine Feed
MEDICATED
Active Drug Ingredient:
OXYTETRACYCLINE**

**This feed contains ____ grams/ton
Oxytetracycline.**

(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. of bodyweight daily.</p> <p>Feed continuously for 7 to 14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to Oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.</p>
<p>10 mg/lb. of bodyweight daily</p> <p>Feed continuously for 14 days.</p>	<p>For breeding swine for control and treatment of Leptospirosis (reducing incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to Oxytetracycline.</p>

Special Considerations: For use in dry animal feed only. Not for use in liquid feed.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Swine Feed
MEDICATED
Active Drug Ingredient:
OXYTETRACYCLINE**

**This feed contains ____ grams/ton
Oxytetracycline.**

(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. of bodyweight daily.</p> <p>Feed continuously for 7 to 14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to Oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.</p>
<p>10 mg/lb. of bodyweight daily</p> <p>Feed continuously for 14 days.</p>	<p>For breeding swine for control and treatment of Leptospirosis (reducing incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to Oxytetracycline.</p>

Special Considerations: For use in dry animal feed only. Not for use in liquid feed.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Swine Feed
MEDICATED
Active Drug Ingredient:
OXYTETRACYCLINE**

**This feed contains ____ grams/ton
Oxytetracycline.**

(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

<p>10 mg/lb. of bodyweight daily.</p> <p>Feed continuously for 7 to 14 days.</p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to Oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.</p>
<p>10 mg/lb. of bodyweight daily</p> <p>Feed continuously for 14 days.</p>	<p>For breeding swine for control and treatment of Leptospirosis (reducing incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to Oxytetracycline.</p>

Special Considerations: For use in dry animal feed only. Not for use in liquid feed.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.