

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: TYLOSIN and
PYRANTEL TARTRATE.**

**This feed contains _____ grams/ton Tylosin and _____
grams/ton Pyrantel Tartrate**

Check Use Level & Indications for Use:

<p>Tylosin: 100 g/ton for at least 3 weeks, followed by 40 g/ton until swine reach market weight</p> <p>Pyrantel Tartrate: 96 g/ton (0.0106%)</p> <p>Feed as the sole ration in a complete feed</p>	<p>For prevention of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>.</p> <p>Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.</p>
<p>Tylosin: 40-100 g/ton for 2-6 weeks after treatment with Tylosin plus vitamins in drinking water (0.25 grams Tylosin per gallon of drinking water for 3-10 days).</p> <p>Pyrantel Tartrate: 96 g/ton (0.0106%)</p>	<p>For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>.</p> <p>Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.</p>

WARNING: Withdraw 24 hours prior to slaughter.

CAUTION: Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing bentonite.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredients: TYLOSIN and
PYRANTEL TARTRATE.**

**This feed contains _____ grams/ton Tylosin and _____
grams/ton Pyrantel Tartrate**

Check Use Level & Indications for Use:

<p>Tylosin: 100 g/ton for at least 3 weeks, followed by 40 g/ton until swine reach market weight</p> <p>Pyrantel Tartrate: 96 g/ton (0.0106%)</p> <p>Feed as the sole ration in a complete feed</p>	<p>For prevention of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>.</p> <p>Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.</p>
<p>Tylosin: 40-100 g/ton for 2-6 weeks after treatment with Tylosin plus vitamins in drinking water (0.25 grams Tylosin per gallon of drinking water for 3-10 days).</p> <p>Pyrantel Tartrate: 96 g/ton (0.0106%)</p>	<p>For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>.</p> <p>Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections.</p>

WARNING: Withdraw 24 hours prior to slaughter.

CAUTION: Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing bentonite.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.