

**Custom Mix Turkey Feed  
MEDICATED**

**Active Drug Ingredient: CHLORTETRACYCLINE**

**This feed contains \_\_\_\_ grams per ton of Chlortetracycline (CTC).**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

200 g/ton	<b>Turkeys (growing)</b> Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
400 g/ton	<b>Turkeys (growing)</b> Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
25 mg/lb. of bodyweight	<b>Turkeys (growing)</b> Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
400 g/ton	<b>Turkeys (poults not over 4 weeks of age)</b> Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to Chlortetracycline.

**WARNING: Do not feed to turkeys producing eggs for human consumption.**

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 01/17

**Custom Mix Turkey Feed  
MEDICATED**

**Active Drug Ingredient: CHLORTETRACYCLINE**

**This feed contains \_\_\_\_ grams per ton of Chlortetracycline (CTC).**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

200 g/ton	<b>Turkeys (growing)</b> Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
400 g/ton	<b>Turkeys (growing)</b> Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
25 mg/lb. of bodyweight	<b>Turkeys (growing)</b> Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to Chlortetracycline. <b>LIMITS FOR USE: Feed continuously for 7 to 14 days.</b>
400 g/ton	<b>Turkeys (poults not over 4 weeks of age)</b> Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to Chlortetracycline.

**WARNING: Do not feed to turkeys producing eggs for human consumption.**

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 01/17