

**Custom Mix Sheep Feed
MEDICATED**

Active Drug Ingredient: LASALOCID

This feed contains _____ grams per ton Lasalocid.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use
20-30 g/ton (0.0022-0.0033%) Feed continuously in a complete feed (Type C) to provide not less than 15 mg nor more than 70 mg/head/day depending on bodyweight. For sheep maintained in confinement.	For the prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>Eimeria crandallis</i> , <i>Eimeria ovinoidalis</i> (<i>Eimeria ninakohlyakimovae</i>), <i>Eimeria parva</i> and <i>Eimeria intricata</i> . Feed _____ to _____ lb./head/day to provide 15 to 70 mg Lasalocid/head/day.

Caution: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A articles or Type B feeds: Mix thoroughly with grain and /or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration which could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration which could be fatal to cattle and sheep.

UK Reg. Services – 11/15

**Custom Mix Sheep Feed
MEDICATED**

Active Drug Ingredient: LASALOCID

This feed contains _____ grams per ton Lasalocid.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use
20-30 g/ton (0.0022-0.0033%) Feed continuously in a complete feed (Type C) to provide not less than 15 mg nor more than 70 mg/head/day depending on bodyweight. For sheep maintained in confinement.	For the prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>Eimeria crandallis</i> , <i>Eimeria ovinoidalis</i> (<i>Eimeria ninakohlyakimovae</i>), <i>Eimeria parva</i> and <i>Eimeria intricata</i> . Feed _____ to _____ lb./head/day to provide 15 to 70 mg Lasalocid/head/day.

Caution: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A articles or Type B feeds: Mix thoroughly with grain and /or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration which could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration which could be fatal to cattle and sheep.

UK Reg. Services – 11/15

**Custom Mix Sheep Feed
MEDICATED**

Active Drug Ingredient: LASALOCID

This feed contains _____ grams per ton Lasalocid.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use
20-30 g/ton (0.0022-0.0033%) Feed continuously in a complete feed (Type C) to provide not less than 15 mg nor more than 70 mg/head/day depending on bodyweight. For sheep maintained in confinement.	For the prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>Eimeria crandallis</i> , <i>Eimeria ovinoidalis</i> (<i>Eimeria ninakohlyakimovae</i>), <i>Eimeria parva</i> and <i>Eimeria intricata</i> . Feed _____ to _____ lb./head/day to provide 15 to 70 mg Lasalocid/head/day.

Caution: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A articles or Type B feeds: Mix thoroughly with grain and /or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration which could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration which could be fatal to cattle and sheep.

UK Reg. Services – 11/15

**Custom Mix Sheep Feed
MEDICATED**

Active Drug Ingredient: LASALOCID

This feed contains _____ grams per ton Lasalocid.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use
20-30 g/ton (0.0022-0.0033%) Feed continuously in a complete feed (Type C) to provide not less than 15 mg nor more than 70 mg/head/day depending on bodyweight. For sheep maintained in confinement.	For the prevention of coccidiosis caused by <i>Eimeria ovina</i> , <i>Eimeria crandallis</i> , <i>Eimeria ovinoidalis</i> (<i>Eimeria ninakohlyakimovae</i>), <i>Eimeria parva</i> and <i>Eimeria intricata</i> . Feed _____ to _____ lb./head/day to provide 15 to 70 mg Lasalocid/head/day.

Caution: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A articles or Type B feeds: Mix thoroughly with grain and /or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration which could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration which could be fatal to cattle and sheep.

UK Reg. Services – 11/15