

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredients: DECOQUINATE AND MONENSIN

This feed contains _____ grams per ton of Decoquinat

This feed contains _____ grams per ton of Monensin
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

| Use Level | Indications for Use: |
|---|---|
| Decoquinat 12.9 - 90.8 grams/ton | For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle. |
| Monensin 5 – 30 grams/ton | For improved feed efficiency in cattle fed in confinement for slaughter. |
| Feed only to cattle fed in confinement for slaughter. | Feed _____ lb / 100 lb bodyweight/day to provide 22.7 mg Decoquinat/100 lb. bodyweight/day and 50-360mgMonensin/head/day. |

For a feed intake of 2 lb/100 lb. bodyweight, mix 22.7 grams of Decoquinat/ton of feed to provide 22.7 mg/100 lb. bodyweight/day.

Limitations for Use: Feed continuously as the sole ration to provide 22.7 mg Decoquinat per 100 lb. of bodyweight (0.5 mg/kg) per day and 50-360 mg Monensin per head per day. Feed for at least 28 days during periods of coccidiosis exposure or when it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinat. Type C cattle feeds may be manufactured from Decoquinat liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.

Limitations for Liquid Feeds: Liquid Type B feeds must be maintained at pH 5.0-6.5. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily as described, even when not in use. Liquid Type B feeds shall bear an expiration date of 8 weeks after its date of manufacture.

Warning: Do not feed to animals producing milk for food. Do not feed to lactating dairy cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Caution: Do not use Decoquinat in feeds containing bentonite. Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result.

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredients: DECOQUINATE AND MONENSIN

This feed contains _____ grams per ton of Decoquinat

This feed contains _____ grams per ton of Monensin
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

| Use Level | Indications for Use: |
|---|---|
| Decoquinat 12.9 - 90.8 grams/ton | For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle. |
| Monensin 5 – 30 grams/ton | For improved feed efficiency in cattle fed in confinement for slaughter. |
| Feed only to cattle fed in confinement for slaughter. | Feed _____ lb / 100 lb bodyweight/day to provide 22.7 mg Decoquinat/100 lb. bodyweight/day and 50-360mgMonensin/head/day. |

For a feed intake of 2 lb/100 lb. bodyweight, mix 22.7 grams of Decoquinat/ton of feed to provide 22.7 mg/100 lb. bodyweight/day.

Limitations for Use: Feed continuously as the sole ration to provide 22.7 mg Decoquinat per 100 lb. of bodyweight (0.5 mg/kg) per day and 50-360 mg Monensin per head per day. Feed for at least 28 days during periods of coccidiosis exposure or when it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinat. Type C cattle feeds may be manufactured from Decoquinat liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.

Limitations for Liquid Feeds: Liquid Type B feeds must be maintained at pH 5.0-6.5. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily as described, even when not in use. Liquid Type B feeds shall bear an expiration date of 8 weeks after its date of manufacture.

Warning: Do not feed to animals producing milk for food. Do not feed to lactating dairy cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Caution: Do not use Decoquinat in feeds containing bentonite. Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result.