

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredients:

LASALOCID & METHOPRENE

This feed contains _____ grams per ton Lasalocid and _____ % (_____ mg/lb) (S)-Methoprene
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

Lasalocid	
10 to 30 g/ton (0.0011-0.0033%). Feed continuously in complete feed (Type C) to provide not less than 100 mg nor more than 360 mg of Lasalocid/head/day.	For improved feed efficiency in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 100 to 360 mg Lasalocid/head/day.
25 to 30 g/ton (0.0027-0.0033%). Feed continuously in complete feed (Type C) to provide not less than 250 mg nor more than 360 mg of Lasalocid/head/day.	For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter. Feed _____ to _____ lb./head/day to provide 250 to 360 mg Lasalocid/head/day.
60 to 300 mg/head/day. Feed continuously at a rate of no less than 60 mg nor more than 300 mg of Lasalocid/head/day. Hand fed: The medication must be contained in at least one pound of feed. Free Choice: The feed must be manufactured under a feed mill license utilizing an FDA- approved formulation, unless manufactured according to formulas published at 21 CFR 558.311(e)(2), (3) or (4).	For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). Feed _____ to _____ lb./head/day to provide 60 to 300 mg Lasalocid/head/day. Intakes of Lasalocid in excess of 200 mg/head/day have not been shown to more effective than 200mg/head/day.
1 mg/2.2 lb. (1.0 kg) of body weight/day Hand feed to cattle weighing up to 800 lb. with a maximum of 360 mg Lasalocid/head/day.	For control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle. Feed at the rate of _____ lb. per 100 lb. of bodyweight/day to provide 1 mg Lasalocid/2.2 lb. (1.0 kg) of bodyweight/day. (Up to a maximum of 360 mg/head/day)
(S)-Methoprene	
Feed 22.7-45.4 mg /100 lb bodyweight/month.	Insect growth regulator for continuous feeding during the fly season to prevent the development of horn flies in the manure of treated cattle.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. For Type A articles and Type B feeds (cattle and sheep): Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess concentration of Lasalocid that could be fatal to cattle and sheep. In addition, inadequate mixing (recirculation or agitation) of liquid Type B feed may result in an excess Lasalocid concentration that could be fatal to cattle and sheep.

Caution: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

Special Considerations: Type C cattle feeds may be manufactured from Lasalocid liquid Type B feeds which have a pH of 4.0 to 8.0 and bear appropriate mixing directions as follows: 1) For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. 2) For liquid Type B feeds stored in mechanical, air, or other agitation type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.