

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains ____ grams per ton of MONENSIN.
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

	Continuously as sole ration 90-110 g/ton	Chickens (broilers) As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>Eimeria tenella</i> , <i>Eimeria acervulina</i> , <i>Eimeria brunetti</i> , <i>Eimeria mivati</i> and <i>Eimeria maxima</i> . WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains. LIMITATIONS FOR USE: Feed continuously as the sole ration.
	90-110 g/ton	Chickens (replacements, intended for use as caged layers) As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>Eimeria tenella</i> , <i>Eimeria acervulina</i> , <i>Eimeria brunetti</i> , <i>Eimeria mivati</i> and <i>Eimeria maxima</i> . WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: MONENSIN

This feed contains ____ grams per ton of MONENSIN.
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

	Continuously as sole ration 90-110 g/ton	Chickens (broilers) As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>Eimeria tenella</i> , <i>Eimeria acervulina</i> , <i>Eimeria brunetti</i> , <i>Eimeria mivati</i> and <i>Eimeria maxima</i> . WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains. LIMITATIONS FOR USE: Feed continuously as the sole ration.
	90-110 g/ton	Chickens (replacements, intended for use as caged layers) As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>Eimeria tenella</i> , <i>Eimeria acervulina</i> , <i>Eimeria brunetti</i> , <i>Eimeria mivati</i> and <i>Eimeria maxima</i> . WARNING: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.