

**Custom Mix Dairy Cattle Feed  
TYPE C MEDICATED**

**Active Drug Ingredient: MONENSIN**

This feed contains \_\_\_\_ grams per ton Monensin.  
Grams per ton divided by 2 is equivalent to milligrams per pound.

**Use Level for TMR herds**

**Indications for Use**

<b>Feed ____ to ____ lbs/hd/day to provide 11 to 22 g/ton on a 100% dry matter basis.</b> Feed continuously to dry and lactating dairy cows a total mixed ration (complete feed) containing 11 to 22 g/ton Monensin on a 100% dry matter basis	For increased milk production efficiency (production of marketable solids – corrected milk per unit of feed intake) in dairy cows.
---	--

**Caution:** Do not allow horses or other equine access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**You May Notice:**

- Reduced voluntary feed intake in dairy cows fed Monensin. This reduction increases with higher doses of Monensin fed. Rule out Monensin as the cause of reduced feed intake before attributing to other causes such as illness, feed management, or the environment.
- Reduced milk fat percentage in dairy cows fed Monensin. This reduction increases with higher doses of Monensin fed.
- Increased incidence and treatment of cystic ovaries and metritis in dairy cows fed Monensin.
- Reduced conception rates, increased services per animal, and extended days open and corresponding calving intervals in dairy cows fed Monensin.

Have a comprehensive and ongoing nutritional, reproductive, and heard health program in place when feeding Monensin to dairy cows.

**Warning:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**Custom Mix Dairy Cattle Feed  
TYPE C MEDICATED**

**Active Drug Ingredient: MONENSIN**

This feed contains \_\_\_\_ grams per ton Monensin.  
Grams per ton divided by 2 is equivalent to milligrams per pound.

**Use Level for TMR herds**

**Indications for Use**

<b>Feed ____ to ____ lbs/hd/day to provide 11 to 22 g/ton on a 100% dry matter basis.</b> Feed continuously to dry and lactating dairy cows a total mixed ration (complete feed) containing 11 to 22 g/ton Monensin on a 100% dry matter basis	For increased milk production efficiency (production of marketable solids – corrected milk per unit of feed intake) in dairy cows.
---	--

**Caution:** Do not allow horses or other equine access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**You May Notice:**

- Reduced voluntary feed intake in dairy cows fed Monensin. This reduction increases with higher doses of Monensin fed. Rule out Monensin as the cause of reduced feed intake before attributing to other causes such as illness, feed management, or the environment.
- Reduced milk fat percentage in dairy cows fed Monensin. This reduction increases with higher doses of Monensin fed.
- Increased incidence and treatment of cystic ovaries and metritis in dairy cows fed Monensin.
- Reduced conception rates, increased services per animal, and extended days open and corresponding calving intervals in dairy cows fed Monensin.

Have a comprehensive and ongoing nutritional, reproductive, and heard health program in place when feeding Monensin to dairy cows.

**Warning:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.