

Custom Mix Goat Feed
MEDICATED

Active Drug Ingredient: Morantel Tartrate

This feed contains _____ grams per ton Morantel Tartrate

0.44 g/100 lb bodyweight	For removal and control of mature gastrointestinal nematode infections of goats including <i>Haemonchus contortus</i> , <i>Ostertagia (Teladorsagia) circumcincta</i> and <i>Trichostrongylus axei</i> . Feed _____ lb. per 100 lb of bodyweight to provide 0.44 g Morantel Tartrate/ 100 lb bodyweight.
--------------------------	---

WARNING: Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats.

CAUTION: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

LIMITATIONS FOR USE: Fresh water should be available at all times. When medicated feed is consumed resume normal feeding. Conditions of constant worm exposure may require retreatment in 2-4 weeks.

UK Regulatory Services – 12/20

Custom Mix Goat Feed
MEDICATED

Active Drug Ingredient: Morantel Tartrate

This feed contains _____ grams per ton Morantel Tartrate

0.44 g/100 lb bodyweight	For removal and control of mature gastrointestinal nematode infections of goats including <i>Haemonchus contortus</i> , <i>Ostertagia (Teladorsagia) circumcincta</i> and <i>Trichostrongylus axei</i> . Feed _____ lb. per 100 lb of bodyweight to provide 0.44 g Morantel Tartrate/ 100 lb bodyweight.
--------------------------	---

WARNING: Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats.

CAUTION: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

LIMITATIONS FOR USE: Fresh water should be available at all times. When medicated feed is consumed resume normal feeding. Conditions of constant worm exposure may require retreatment in 2-4 weeks.

UK Regulatory Services – 12/20

Custom Mix Goat Feed
MEDICATED

Active Drug Ingredient: Morantel Tartrate

This feed contains _____ grams per ton Morantel Tartrate

0.44 g/100 lb bodyweight	For removal and control of mature gastrointestinal nematode infections of goats including <i>Haemonchus contortus</i> , <i>Ostertagia (Teladorsagia) circumcincta</i> and <i>Trichostrongylus axei</i> . Feed _____ lb. per 100 lb of bodyweight to provide 0.44 g Morantel Tartrate/ 100 lb bodyweight.
--------------------------	---

WARNING: Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats.

CAUTION: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

LIMITATIONS FOR USE: Fresh water should be available at all times. When medicated feed is consumed resume normal feeding. Conditions of constant worm exposure may require retreatment in 2-4 weeks.

UK Regulatory Services – 12/20

Custom Mix Goat Feed
MEDICATED

Active Drug Ingredient: Morantel Tartrate

This feed contains _____ grams per ton Morantel Tartrate

0.44 g/100 lb bodyweight	For removal and control of mature gastrointestinal nematode infections of goats including <i>Haemonchus contortus</i> , <i>Ostertagia (Teladorsagia) circumcincta</i> and <i>Trichostrongylus axei</i> . Feed _____ lb. per 100 lb of bodyweight to provide 0.44 g Morantel Tartrate/ 100 lb bodyweight.
--------------------------	---

WARNING: Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats.

CAUTION: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

LIMITATIONS FOR USE: Fresh water should be available at all times. When medicated feed is consumed resume normal feeding. Conditions of constant worm exposure may require retreatment in 2-4 weeks.

UK Regulatory Services – 12/20