

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20

Custom Mix Growing Pheasant Feed MEDICATED

Active Drug Ingredient: AMPROLIUM

This feed contains 159 grams per ton of Amprolium.

Use Level	Indications for Use
159 g/ton (0.175%)	For the prevention of coccidiosis in growing pheasants caused by <i>Eimeria colchici</i> , <i>Eimeria duodenalis</i> and <i>Eimeria phasiani</i> .

WARNING: Use as sole source of Amprolium. Fertility, hatchability and other reproductive data are not available on Amprolium in breeding pheasants.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS FOR USE: Feed continuously as sole ration.

UK Regulatory Services – 12/20