

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20

**Custom Mix Quail Feed
MEDICATED**

Active Drug Ingredient: **Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

5 to 20 g/ton.	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.
200 g/ton. Limitations for Use: Feed continuously as the sole ration to growing quail.	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to BMD.

UK Regulatory Services -12/20