Custom Mix Sheep Feed MEDICATED

Active Drug Ingredient: DECOQUINATE

This feed contains	grams per ton decoquinate.
(Grams per ton divided by	y 2 is equivalent to milligrams per pound.

Check Use Level	Indications for Use
For Young Sheep	For the prevention of coccidiosis in young sheep caused by Eimeria bakuensis, Eimeria crandallis,
12.9 – 90.8 g/ton.	Eimeria ovinoidalis and Eimeria parva.
Limitations for use:	Feed lb./100 lb. of bodyweight/day to
Feed Type C medicated	provide 22.7 mg/100 lb. (0.5mg/kg) of
feed or milk replacer at	bodyweight/day.
a rate to provide 22.7	
mg/100 lb (0.5 mg/kg) of	
bodyweight per day.	
For Young Sheep	For the prevention of coccidiosis caused by Eimeria bakuensis, Eimeria crandallis, Eimeria ovinodialis
90.9 – 535.7 g/ton.	and Eimeria parva.
Limitations for use:	
Feed Type C medicated	Feed lb./100 lb. of bodyweight/day to
feed supplements as a	provide 22.7 mg/100 lb. (0.5mg/kg) of
top dress or mix into the	bodyweight/day.
daily ration to provide	
22.7 mg/100 lb (0.5	
mg/kg) of	
bodyweight/day.	

WARNING: Do not feed to sheep producing milk for human consumption.

CAUTION: Do not use Decoquinate in feeds containing bentonite.

<u>LIMITATIONS FOR USE</u>: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/19

Custom Mix Sheep Feed MEDICATED

Active Drug Ingredient: DECOQUINATE

•	This feed contains	grams per ton decoquinate.
(Grams per ton divided by	2 is equivalent to milligrams per pound

Check Use Level	Indications for Use
For Young Sheep	For the prevention of coccidiosis in young sheep
12.9 – 90.8 g/ton.	caused by Eimeria bakuensis, Eimeria crandallis, Eimeria ovinoidalis and Eimeria parva.
Limitations for use:	Feed lb./100 lb. of bodyweight/day to
Feed Type C medicated	provide 22.7 mg/100 lb. (0.5mg/kg) of
feed or milk replacer at	bodyweight/day.
a rate to provide 22.7	
mg/100 lb (0.5 mg/kg) of	
bodyweight per day.	
T. W. GI	
For Young Sheep	For the prevention of coccidiosis caused by <i>Eimeria</i> bakuensis, Eimeria crandallis, Eimeria ovinodialis
90.9 – 535.7 g/ton.	and Eimeria parva.
Limitations for use:	
Feed Type C medicated	Feedlb./100 lb. of bodyweight/day to
feed supplements as a	provide 22.7 mg/100 lb. (0.5mg/kg) of
top dress or mix into the	bodyweight/day.
daily ration to provide	
22.7 mg/100 lb (0.5	
mg/kg) of	
bodyweight/day.	

WARNING: Do not feed to sheep producing milk for human consumption.

<u>CAUTION:</u> Do not use Decoquinate in feeds containing bentonite.

<u>LIMITATIONS FOR USE</u>: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid Type B feed.

UK Regulatory Services – 01/19