

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredient: Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

10 to 30 g/ton. For growing and finishing swine.	For increased rate of weight gain and improved feed efficiency.
250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration. Not for use in swine weighing more than 250 lb.	For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.
250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration.	For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours.

UK Regulatory Services – 12/20

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredient: Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

10 to 30 g/ton. For growing and finishing swine.	For increased rate of weight gain and improved feed efficiency.
250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration. Not for use in swine weighing more than 250 lb.	For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.
250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration.	For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours.

UK Regulatory Services – 12/20

**Custom Mix Swine Feed
MEDICATED**

**Active Drug Ingredient: Bacitracin
Methylenedisalicylate (BMD)**

This feed contains ____ grams per ton of BMD.

Check Use Level & Indications for Use:

10 to 30 g/ton. For growing and finishing swine.	For increased rate of weight gain and improved feed efficiency.
250 g/ton. For growing and finishing swine. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration. Not for use in swine weighing more than 250 lb.	For control of swine dysentery (bloody scours) associated with <i>Treponema hyodysenteria</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.
250 g/ton. For pregnant sows. Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylenedisalicylate should be the sole ration.	For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours.

UK Regulatory Services – 12/20