

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).

(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

Calves (up to 250 lb.) 10 mg/lb. bodyweight/day*	For treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to chlortetracycline. Feed ____ lb./100 lb. bodyweight to provide 10 mg/lb. bodyweight/day of chlortetracycline. LIMITS FOR USE: In milk replacers or starter feeds.
Growing Cattle (over 400 lb.) 70 mg/head/day*	Reduction of incidence of liver abscesses. Feed ____ lb./head/day to provide 70 mg chlortetracycline.
Beef cattle and dairy replacement heifers 350 mg/head/day*	Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella spp.</i> susceptible to Chlortetracycline. Feed ____ lb./head/day to provide 350 mg chlortetracycline.
Beef cattle (under 700 lb.) 350 mg/head/day*	For control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. Feed ____ lb./head/day to provide 350 mg chlortetracycline.
Beef cattle (over 700 lb.) 0.5 mg/lb. bodyweight/day*	Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. Feed ____ lb. daily to provide 0.5 mg of chlortetracycline/lb. bodyweight. <i>Example: To provide 0.5 mg/lb. bodyweight at a feed intake of 0.5 lb/100 lb. bodyweight, mix a feed containing 200g Chlortetracycline</i>
Calves, beef and non-lactating dairy cattle 10 mg/lb. bodyweight/day*	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. Feed ____ lb./100 lb. bodyweight/day to provide 10 mg chlortetracycline/lb. bodyweight/day. Limitations for Use: Feed for not more than 5 days. Feed approximately 400 g/ton varying with bodyweight and feed consumption to provide 10 mg/lb. per day of chlortetracycline.
Beef and non-lactating dairy cattle 0.5-2.0 mg/lb. bodyweight/day	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline. LIMITATIONS: In free-choice cattle feeds such as feed blocks or salt-mineral mixes; Free choice feed must be manufactured from NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin) using an FDA-approved formulation. Feed mill license not required if following free-choice loose-mineral formulation published in 21 CFR 558.128.
Calves, beef and non-lactating dairy cattle 500 – 4,000 g/ton*	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. LIMITATIONS: Feed continuously for not more than 5 days to provide 10 mg/lb. bodyweight per day; must be manufactured under NADA 048-761 (Aureomycin) or ANADA 200-510 (Deracin).
Beef and non-lactating dairy cattle 6,000 g/ton	As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to Chlortetracycline.
Calves, beef and non-lactating dairy cattle 4,000-20,000 g/ton	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline. LIMITATIONS: As a top dress, varying with bodyweight and feed consumption, to provide 10 mg/lb. bodyweight per day. Treat for not more than 5 days, must be manufactured from Aureomycin Type A medicated articles manufactured under NADA 048-761 or Deracin Type A medicated articles manufactured under ANADA 200-510.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.

*Withdrawal times may vary by manufacturer. Read label for specific withdrawal times.

Withdrawal time for the use checked above is _____ prior to slaughter.

CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.