

**Custom Mix Cattle Feed  
Chlortetracycline / Decoquinatone  
MEDICATED**

**Active Drug Ingredients:**

\_\_\_\_\_ grams per ton **CHLORTETRACYCLINE (CTC)** and

\_\_\_\_\_ grams per ton **DECOQUINATE**  
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Use Level	Indications for Use:
<b>CTC</b> <b>500 – 4000 g/ton</b>  and  <b>Decoquinatone</b> <b>12.9-90.8 g/ton</b>	<b>Calves, beef and nonlactating dairy cattle</b> For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> . For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.  <b>Feed _____ lbs./100 lb bodyweight/day to provide 22.7 mg Decoquinatone and 1 g CTC per 100 lb. bodyweight per day for not more than 5 days.</b>
<b>CTC</b> <b>4,000 to 20,000 g/ton</b>  and  <b>Decoquinatone</b> <b>90.9 to 535.7 g/ton</b>	<b>Calves, beef and nonlactating dairy cattle</b> For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> . For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline.  <b>LIMITATIONS FOR USE: Feed Type C medicated supplements as a top dress or mix into daily ration to provide 22.7 mg Decoquinatone and 1 g CTC per 100 lb. bodyweight per day for not more than 5 days.</b>

**LIMITATIONS FOR USE:** Feed Type C feed at a rate to provide 22.7 mg Decoquinatone and 1 g Chlortetracycline per 100 lb. of bodyweight (0.5 mg/kg) per day for not more than 5 days. When consumed, feed 22.7 mg Decoquinatone per 100 lb. bodyweight per day for total of 28 days to prevent coccidiosis. Type C feed may be prepared from dry or liquid Type B feed containing 0.0125-0.5% Decoquinatone. Type C cattle feeds may be manufactured from Decoquinatone liquid Type B feeds having a pH between 5.0 and 6.5 and containing a suspending agent to maintain a viscosity of not less than 500 centipoises.

**WARNING:** Do not feed to animals producing milk for food. A withdrawal period has not been established for this product in pre-ruminating calves. Do not feed to calves to be processed for veal. Withdraw 24 hours prior to slaughter when manufactured from ChlorMax Type A medicated articles and Pennchlor Type A medicated articles. Zero withdrawal time when manufactured from Aureomycin Type A medicated articles manufactured by NADA sponsor No. 054771 (Zoetis) and Deracin Type A medicated articles manufactured by ANADA sponsor No. 069254 (Pharmgate Animal Health).

**CAUTION:** Do not use Decoquinatone in feeds containing bentonite.

**CAUTION:** Federal law restricts medicated feed containing this Veterinary feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.