

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

10 mg/lb. bodyweight daily* Feed continuously daily for 7 to 14 days. Note: Calf weighing 100 lb consuming 1 lb. of dry milk replacer mixed with 1 gal. water.	Calves (milk replacers): Treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to Oxytetracycline.
75 mg/head/day For Beef cattle. Use continuously	For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses. Feed _____ lb./head/day.
0.5 to 2.0 g/head/day For beef cattle. Feed 3-5 days before and after arrival in feedlot.	Prevention and treatment of the early stages of shipping fever complex. Feed __ to __ lb./head/day.
10 mg/lb. bodyweight daily For calves, beef cattle and nonlacting dairy cattle. Feed continuously for 7-14 days.	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. Feed _____ lb./100 lb. bodyweight/day.

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Cattle Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains _____ grams/ton Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound.)

Check Use Level & Indications for Use:

10 mg/lb. bodyweight daily* Feed continuously daily for 7 to 14 days. Note: Calf weighing 100 lb consuming 1 lb. of dry milk replacer mixed with 1 gal. water.	Calves (milk replacers): Treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to Oxytetracycline.
75 mg/head/day For Beef cattle. Use continuously	For growing cattle (over 400 lb.) for reduction of liver condemnation due to liver abscesses. Feed _____ lb./head/day.
0.5 to 2.0 g/head/day For beef cattle. Feed 3-5 days before and after arrival in feedlot.	Prevention and treatment of the early stages of shipping fever complex. Feed __ to __ lb./head/day.
10 mg/lb. bodyweight daily For calves, beef cattle and nonlacting dairy cattle. Feed continuously for 7-14 days.	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. Feed _____ lb./100 lb. bodyweight/day.

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements.

WARNING: Withdrawal times may vary by manufacturer. Read label for specific withdrawal time. When used in milk replacers, the treatment claim (10mg/lb.) is limited to bacterial enteritis caused by *Escherichia coli* only.

LIMITATIONS FOR USE: For product(s) manufactured by NADA sponsor No. 069254 (Pharmgate Animal Health), withdraw 5 days before slaughter when fed at 10 mg/lb. bodyweight daily. For product(s) manufactured by NADA sponsor No. 066104 (Phibro Animal Health), zero-day withdrawal.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.