## Custom Mix Cattle Feed MEDICATED

**Active Drug Ingredient: TYLOSIN** 

This feed contains \_\_\_\_ grams per ton tylosin. (Grams per ton divided by 2 is equivalent to milligrams per pound.)

For beef cattle	For reduction of incidence of
	liver abscesses in beef cattle
8 to 10 g/ton* to	caused by Fusobacterium
provide 60-90 mg	necrophorum and
/head/day (Feed	Arcanobacterium
continuously.)	(Actinomyces) pyogenes.
*90% dry matter basis.	Feed to lbs/head/day continuously to
	provide 60-90 mg
	Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 1/19

## Custom Mix Cattle Feed MEDICATED

**Active Drug Ingredient: TYLOSIN** 

This feed contains grams per ton tylosin. (Grams per ton divided by 2 is equivalent to milligrams per pound.)

For beef cattle	For reduction of incidence of
	liver abscesses in beef cattle
8 to 10 g/ton* to	caused by Fusobacterium
provide 60-90 mg	necrophorum and
/head/day (Feed	Arcanobacterium
continuously.)	(Actinomyces) pyogenes.
*90% dry matter basis.	Feed to lbs/head/day continuously to provide 60-90 mg
	Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**UK Regulatory Services – 1/19** 

## Custom Mix Cattle Feed MEDICATED

**Active Drug Ingredient: TYLOSIN** 

This feed contains \_\_\_\_ grams per ton tylosin. (Grams per ton divided by 2 is equivalent to milligrams per pound.)

For beef cattle 8 to 10 g/ton* to provide 60-90 mg /head/day (Feed	For reduction of incidence of liver abscesses in beef cattle caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium</i>
continuously.)	(Actinomyces) pyogenes.
*90% dry matter basis.	Feed to lbs/head/day continuously to provide 60-90 mg Tylosin/head/day.

Notice: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

LIMITATIONS FOR LIQUID FEEDS: Liquid Type B feeds must be maintained at ph 4.5-6.0. Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate or agitate daily, even when not in use. Liquid Type B feeds shall bear an expiration date of 31 days after its date of manufacture.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**UK Regulatory Services – 1/19**