

**Custom Mix Chicken Feed  
MEDICATED**

**Active Drug Ingredient: CHLORTETRACYCLINE**

**This feed contains \_\_\_\_ grams per ton of Chlortetracycline (CTC).**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

<b>100-200 g/ton</b>	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.</b>
<b>200-400 g/ton</b>	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.</b>
<b>500 g/ton</b>	Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: (1) Feed for 5 days. Zero-day withdrawal time when formulated from Aureomycin Type A medicated articles or Type B medicated feeds under NADA 48-761 or Deracin Type A medicated articles or Type B medicated feeds under ANADA 200-510. (2) All other Chlortetracycline premixes: Feed for 5 days. Withdraw 24 hours prior to slaughter. Do not feed to chickens producing eggs for human consumption.</b>

**WARNING:** Varies by manufacturer. Refer to specific product label.

**VFD CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Custom Mix Chicken Feed  
MEDICATED**

**Active Drug Ingredient: CHLORTETRACYCLINE**

**This feed contains \_\_\_\_ grams per ton of Chlortetracycline (CTC).**  
(Grams per ton divided by 2 is equivalent to milligrams per pound)

**Check Use Level & Indications for Use:**

<b>100-200 g/ton</b>	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.</b>
<b>200-400 g/ton</b>	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: Feed continuously for 7 to 14 days. For all Chlortetracycline products except Aureomycin or Deracin: Do not feed to chickens producing eggs for human consumption.</b>
<b>500 g/ton</b>	Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to Chlortetracycline. <b>LIMITATIONS FOR USE: (1) Feed for 5 days. Zero-day withdrawal time when formulated from Aureomycin Type A medicated articles or Type B medicated feeds under NADA 48-761 or Deracin Type A medicated articles or Type B medicated feeds under ANADA 200-510. (2) All other Chlortetracycline premixes: Feed for 5 days. Withdraw 24 hours prior to slaughter. Do not feed to chickens producing eggs for human consumption.</b>

**WARNING:** Varies by manufacturer. Refer to specific product label.

**VFD CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.