

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains ____ grams per ton of Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

100-200 g/ton Feed continuously for 7-14 days	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
400 g/ton Feed continuously for 7-14 days	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Oxytetracycline.
500 g/ton Feed continuously for 5 days	Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to Oxytetracycline.

WARNING: At 500 g/ton use level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower use levels. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues. Do not administer to chickens producing eggs for human consumption.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 12/19

**Custom Mix Chicken Feed
MEDICATED**

Active Drug Ingredient: OXYTETRACYCLINE

This feed contains ____ grams per ton of Oxytetracycline.
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

100-200 g/ton Feed continuously for 7-14 days	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline.
400 g/ton Feed continuously for 7-14 days	Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to Oxytetracycline.
500 g/ton Feed continuously for 5 days	Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to Oxytetracycline.

WARNING: At 500 g/ton use level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower use levels. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues. Do not administer to chickens producing eggs for human consumption.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 12/19