

**Custom Mix Sheep Feed
MEDICATED**

**Active Drug Ingredient:
OXYTETRACYCLINE**

This feed contains ____ grams per ton of
Oxytetracycline.

(Grams per ton divided by 2 is equivalent to
milligrams per pound.)

Indications for Use:

| | |
|---|--|
| 10 mg/lb. of bodyweight daily. | Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. |
| Feed continuously for 7-14 days. | Feed ____ lbs of this feed/100 lbs of bodyweight/day to provide 10 mg/lb of bodyweight/day of Oxytetracycline. |

Warning: At 10 mg/lb dosage, 5 day withdrawal required.

SPECIAL CONSIDERATION: For use in dry animal feed
only. Not for use in liquid feed supplements.

VFD CAUTION: Federal law restricts medicated feed
containing this Veterinary Feed Directive (VFD) drug to use
by or on the order of a licensed veterinarian.

UK Regulatory Services – 1/19

**Custom Mix Sheep Feed
MEDICATED**

**Active Drug Ingredient:
OXYTETRACYCLINE**

This feed contains ____ grams per ton of
Oxytetracycline.

(Grams per ton divided by 2 is equivalent to
milligrams per pound.)

Indications for Use:

| | |
|---|--|
| 10 mg/lb. of bodyweight daily. | Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. |
| Feed continuously for 7-14 days. | Feed ____ lbs of this feed/100 lbs of bodyweight/day to provide 10 mg/lb of bodyweight/day of Oxytetracycline. |

Warning: At 10 mg/lb dosage, 5 day withdrawal required.

SPECIAL CONSIDERATION: For use in dry animal feed
only. Not for use in liquid feed supplements.

VFD CAUTION: Federal law restricts medicated feed
containing this Veterinary Feed Directive (VFD) drug to use
by or on the order of a licensed veterinarian.

UK Regulatory Services – 1/19

**Custom Mix Sheep Feed
MEDICATED**

**Active Drug Ingredient:
OXYTETRACYCLINE**

This feed contains ____ grams per ton of
Oxytetracycline.

(Grams per ton divided by 2 is equivalent to
milligrams per pound.)

Indications for Use:

| | |
|---|--|
| 10 mg/lb. of bodyweight daily. | Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. |
| Feed continuously for 7-14 days. | Feed ____ lbs of this feed/100 lbs of bodyweight/day to provide 10 mg/lb of bodyweight/day of Oxytetracycline. |

Warning: At 10 mg/lb dosage, 5 day withdrawal required.

SPECIAL CONSIDERATION: For use in dry animal feed
only. Not for use in liquid feed supplements.

VFD CAUTION: Federal law restricts medicated feed
containing this Veterinary Feed Directive (VFD) drug to use
by or on the order of a licensed veterinarian.

UK Regulatory Services – 1/19

**Custom Mix Sheep Feed
MEDICATED**

**Active Drug Ingredient:
OXYTETRACYCLINE**

This feed contains ____ grams per ton of
Oxytetracycline.

(Grams per ton divided by 2 is equivalent to
milligrams per pound.)

Indications for Use:

| | |
|---|--|
| 10 mg/lb. of bodyweight daily. | Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline. |
| Feed continuously for 7-14 days. | Feed ____ lbs of this feed/100 lbs of bodyweight/day to provide 10 mg/lb of bodyweight/day of Oxytetracycline. |

Warning: At 10 mg/lb dosage, 5 day withdrawal required.

SPECIAL CONSIDERATION: For use in dry animal feed
only. Not for use in liquid feed supplements.

VFD CAUTION: Federal law restricts medicated feed
containing this Veterinary Feed Directive (VFD) drug to use
by or on the order of a licensed veterinarian.

UK Regulatory Services – 1/19