

**Custom Mix Swine Feed  
MEDICATED**  
**Active Drug Ingredients:**  
**NEOMYCIN/OXYTETRACYCLINE**

**This feed contains \_\_\_\_\_ grams/ton  
Neomycin/Oxytetracycline.**  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b><u>Swine</u></b>	For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>Escherichia coli</i> susceptible to Neomycin.
To provide 10 mg/lb. of bodyweight daily.  Limitations for use: Feed continuously for 7 to 14 days. Withdraw 5 days before slaughter.	
<b><u>Breeding Swine</u></b>	For control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to Oxytetracycline.
To provide 10 mg/lb. of bodyweight daily  Limitations for use: Feed continuously for not more than 14 days. Withdraw 5 days before slaughter.	

**VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.**

UK Reg. Services – 01/19

**Custom Mix Swine Feed  
MEDICATED**  
**Active Drug Ingredients:**  
**NEOMYCIN/OXYTETRACYCLINE**

**This feed contains \_\_\_\_\_ grams/ton  
Neomycin/Oxytetracycline.**  
(Grams per ton divided by 2 is equivalent to  
milligrams per pound.)

**Check Use Level & Indications for Use:**

<b><u>Swine</u></b>	For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>Escherichia coli</i> susceptible to Neomycin.
To provide 10 mg/lb. of bodyweight daily.  Limitations for use: Feed continuously for 7 to 14 days. Withdraw 5 days before slaughter.	
<b><u>Breeding Swine</u></b>	For control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to Oxytetracycline.
To provide 10 mg/lb. of bodyweight daily  Limitations for use: Feed continuously for not more than 14 days. Withdraw 5 days before slaughter.	

**VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.**

UK Reg. Services – 01/19