

**Custom Mix Swine Feed  
MEDICATED**

**Active Drug Ingredients: TYLOSIN**

**This feed contains \_\_\_\_\_ grams per ton tylosin.**

Check Use Level & Indications for Use:

<b>100 g/ton for 21 days.</b>	For control of swine dysentery associated with <i>Brachyspira hyodysenteriae</i> ; and for control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .
<b>100 g/ton for at least 3 weeks</b>  Or  <b>40 g/ton (Include 100 g/ton of Tylosin for at least 3 weeks, followed by 40 g/ton until market weight.)</b>	For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i> ; and for control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .
<b>100 g/ton</b>	Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis.
<b>40-100 g/ton for 2-6 weeks immediately after treatment with Tylosin Tartrate (250 mg Tylosin per gallon in drinking water for 3-10 days)</b>	Treatment and control of swine dysentery associated with <i>Brachyspira hyodysenteriae</i> . For control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .

**Notice:** Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 12/17

**Custom Mix Swine Feed  
MEDICATED**

**Active Drug Ingredients: TYLOSIN**

**This feed contains \_\_\_\_\_ grams per ton tylosin.**

Check Use Level & Indications for Use:

<b>100 g/ton for 21 days.</b>	For control of swine dysentery associated with <i>Brachyspira hyodysenteriae</i> ; and for control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .
<b>100 g/ton for at least 3 weeks</b>  Or  <b>40 g/ton (Include 100 g/ton of Tylosin for at least 3 weeks, followed by 40 g/ton until market weight.)</b>	For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i> ; and for control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .
<b>100 g/ton</b>	Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis.
<b>40-100 g/ton for 2-6 weeks immediately after treatment with Tylosin Tartrate (250 mg Tylosin per gallon in drinking water for 3-10 days)</b>	Treatment and control of swine dysentery associated with <i>Brachyspira hyodysenteriae</i> . For control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> .

**Notice:** Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 12/17