

**Custom Mix Turkey Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

200 g/ton	Turkeys (growing) Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
400 g/ton	Turkeys (growing) Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
25 mg/lb. of bodyweight	Turkeys (growing) Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
400 g/ton	Turkeys (poults not over 4 weeks of age) Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to Chlortetracycline.

WARNING: Do not feed to turkeys producing eggs for human consumption.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 01/19

**Custom Mix Turkey Feed
MEDICATED**

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains ____ grams per ton of Chlortetracycline (CTC).
(Grams per ton divided by 2 is equivalent to milligrams per pound)

Check Use Level & Indications for Use:

200 g/ton	Turkeys (growing) Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
400 g/ton	Turkeys (growing) Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
25 mg/lb. of bodyweight	Turkeys (growing) Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to Chlortetracycline. LIMITS FOR USE: Feed continuously for 7 to 14 days.
400 g/ton	Turkeys (poults not over 4 weeks of age) Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to Chlortetracycline.

WARNING: Do not feed to turkeys producing eggs for human consumption.

VFD CAUTION: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

UK Regulatory Services – 01/19